


available at www.sciencedirect.comjournal homepage: www.elsevier.com/locate/peptides

Structure-activity relationships for in vitro diuretic activity of CAP2b in the housefly

Ronald J. Nachman^{a,*}, Geoffrey M. Coast^b

^a Areawide Pest Management Research, Southern Plains Agricultural Research Center, USDA, 2881 F/B Road, College Station, TX 77845, USA

^b School of Biological and Chemical Sciences, Birkbeck (University of London), London WC1E 7HX, UK

ARTICLE INFO

Article history:

Received 11 September 2006

Received in revised form

15 September 2006

Accepted 15 September 2006

Published on line 8 December 2006

Keywords:

CAPA gene

Periviscerokinin

CAP2b

Perisymphatic organs

Insect neuropeptide

Malpighian tubule

ABSTRACT

A series of truncated and Ala-replacement analogs of the peptide Manse-CAP2b (pELYAFPRV-NH₂) were assayed for diuretic activity on Malpighian tubules of the housefly *Musca domestica* (*M. domestica*). The C-terminal hexapeptide proved to be the active core, the minimum sequence required to retain significant diuretic activity. However, full activity required the C-terminal heptapeptide, which was equipotent with the most active of the native housefly CAP2b peptides. Replacement of Arg⁷ and Val⁸ with Ala led to inactivity and a large 70-fold drop in potency, respectively, indicating that these were critical residues. The Leu² was semicritical, where a six-fold loss in potency was observed. Conversely, the replacement of all other residues with Ala led to much smaller effects on potency and these positions were considered to be noncritical. This structure-activity relationship data can aid in the design of mimetic agonist/antagonist analogs of this diuretic peptide family with enhanced biostability and bioavailability, as tools for arthropod endocrinologists and as potential pest management agents capable of disrupting the water balance in pest flies.

© 2006 Elsevier Inc. All rights reserved.

1. Introduction

It is well established that neuropeptides are key factors in controlling primary urine production in Malpighian tubules (CRF-like peptides, insect kinins, CAPA peptides) and fluid reabsorption from the hindgut (ion transport peptide (ITP)) [1,29]. CAP2b (pELYAFPRVa), a CAPA peptide (also included in the literature as a member of the periviscerokinin (PVK) family), was first identified as a cardioacceleratory factor in the moth *Manduca sexta* [11]. CAPA peptides, including CAP2b, demonstrate diuretic effects on the Malpighian tubules of a number of flies, including the fruit fly *Drosophila melanogaster*, the housefly *Musca domestica* (*M. domestica*), and the stable fly *Stomoxys calcitrans* [1,2–5,9,10,15,18,21,22,24–26,28]. By contrast,

bioassays on the bug *Rhodnius prolixus* with *Manduca* CAP2b [23,29] indicated that native CAPA peptides might reduce secretion by Malpighian tubules and thus show antidiuretic effects. These peptides are putative hormones typical of the neurosecretory system in the abdominal ventral nerve cord (VNC) of insects. The CAPA gene, which encodes for two or more CAP2b/PVKs and a single pyrokinin (PK), is known from a number of holometabolous insects, including *D. melanogaster* [12,13] and *M. sexta* [14]. Expressed in median neurosecretory neurons of abdominal ganglia, these putative peptide hormones may be released into the hemolymph via perisymphatic organs (PSOs) which are segmentally reiterated neurohemal organs of the abdominal ganglia [7,8]. Larval PSOs of cycloraphous Diptera, however, become incorporated into

* Corresponding author. Tel.: +1 979 260 9315; fax: +1 979 260 9377.

E-mail address: nachman@tamu.edu (R.J. Nachman).

0196-9781/\$ – see front matter © 2006 Elsevier Inc. All rights reserved.

doi:10.1016/j.peptides.2006.09.020

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 15 SEP 2006		2. REPORT TYPE		3. DATES COVERED 00-00-2006 to 00-00-2006	
4. TITLE AND SUBTITLE Structure-activity relationships for in vitro diuretic activity of CAP2b in the housefly				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Department of Agriculture,Areawide Pest Management Research,2881 F/B Road,College Station,TX,77845				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT see report					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

the dorsal ganglionic sheath [20,30,31] during metamorphosis. Direct analysis of abdominal dorsal sheath tissues via MALDI-TOF/TOF mass spectrometry led to the identification of CAP2b/PVK peptides native to the housefly *M. domestica* [18,26,32] and three other species of flies [18,19]. The housefly CAP2b/PVK sequences were recently determined to be AGGTSGLYAFPRVa (Musdo-CAP2b/PVK-1) and ASLFNAPRVa (Musdo-CAP2b/PVK-2) [18]. The octapeptide CAP2b shares a common C-terminal heptapeptide sequence (LYAFPRVa) with the most active housefly CAP2b sequence, Musdo-CAP2b-1, revealing the likely reason that the *Manduca* peptide demonstrates relatively potent diuretic activity in the heterologous housefly Malpighian tubule fluid secretion bioassay.

In this study, we use *Manduca* CAP2b as a model of Musdo-CAP2b-1 and evaluate a series of truncated and Ala-replacement CAP2b analogs in a housefly Malpighian tubule fluid secretion bioassay to identify the active core and those residues most critical to activity.

2. Materials and methods

2.1. Insects

Housefly (*M. domestica*) larvae were obtained from a commercial fish bait supplier. They were held in the laboratory until they pupated and were then moved to a constant temperature room (28 °C; 12 L:12 D light:dark cycle). Flies that emerged within 12 h of one another were kept as separate age cohorts and provided with water and sucrose ad libitum. All of the experiments described in this paper used 3–4 day-old adult female flies.

2.2. Peptide synthesis

The CAP2b analogs were synthesized via Fmoc methodology on Rink Amide resin (Novabiochem, San Diego, CA) using Fmoc protected amino acids (Advanced Chemtech, Louisville, KY) on an ABI 433A peptide synthesizer (Applied Biosystems, Foster City, CA) under previously described conditions [17]. Crude products were purified on a Waters C₁₈ Sep Pak cartridge and a Delta Pak C₁₈ reverse-phase column (8 mm × 100 mm, 15 µm particle size, 100 Å pore size) on a Waters 510 HPLC controlled with a Millennium 2010 chromatography manager system (Waters, Milford, MA) with detection at 214 nm at ambient temperature. Solvent A = 0.1% aqueous trifluoroacetic acid (TFA); Solvent B = 80% aqueous acetonitrile containing 0.1% TFA. Conditions: initial solvent consisting of 20% B was followed by the Waters linear program to 100% B over 40 min; flow rate, 2.0 ml/min. Delta-Pak C-18 retention times: Manse-CAP2b[2–8], 9.0 min; Manse-CAP2b[3–8], 7.5 min; Manse-CAP2b[4–8], 5.0 min; Manse-CAP2b[5–8], 4.5 min; Manse-CAP2b[Ala¹], 10.5 min; Manse-CAP2b[Ala²], 9.0 min; Manse-CAP2b[Ala³], 4.5 min; Manse-CAP2b[Ala⁵], 7.5 min; Manse-CAP2b[Ala⁶], 12.5 min; Manse-CAP2b[Ala⁷], 10.5 min; Manse-CAP2b[Ala⁸], 9.0 min. The peptides were further purified on a Waters Protein Pak I125 column (7.8 mm × 300 mm)(Milligen Corp., Milford, MA). Conditions: flow rate: 2.0 ml/min; isocratic with solvent consisting of 80% acetonitrile made to 0.01% TFA. WatPro retention times: Manse-CAP2b[2–8], 7.0 min; Manse-CAP2b[3–8], 8.0 min;

Manse-CAP2b[4–8], 11.5 min; Manse-CAP2b[5–8], 10.5 min; Manse-CAP2b[Ala¹], 15.5 min; Manse-CAP2b[Ala²], 15.0 min; Manse-CAP2b[Ala³], 13.5 min; Manse-CAP2b[Ala⁵], 16.5 min; Manse-CAP2b[Ala⁶], 11.0 min; Manse-CAP2b[Ala⁷], 5.5 min; Manse-CAP2b[Ala⁸], 15.0 min. Amino acid analysis was carried out under previously reported conditions [17] and used to quantify the peptide and to confirm identity, leading to the following analyses: Manse-CAP2b[2–8]: A[1.0], F[1.0], L[1.0], P[1.0], R[1.0], Y[1.0], V[1.0]; Manse-CAP2b[3–8]: A[1.0], F[1.0], P[1.1], R[1.0], Y[1.0], V[1.0]; Manse-CAP2b[4–8]: A[0.8], F[1.0], P[1.0], R[1.0], V[0.9]; Manse-CAP2b[5–8]: F[1.0], P[1.0], R[1.0], V[1.0]; Manse-CAP2b[Ala¹]: A[1.9], F[1.0], L[1.0], P[1.0], R[1.0], Y[1.1], V[0.9]; Manse-CAP2b[Ala²]: A[2.0], E[0.9], F[1.0], P[0.9], R[1.0], Y[1.1], V[1.0]; Manse-CAP2b[Ala³]: A[2.0], E[0.9], F[1.0], L[1.0], P[1.0], R[1.0], V[0.9]; Manse-CAP2b[Ala⁵]: A[2.0], E[0.8], L[1.0], P[0.9], R[1.0], Y[1.2], V[0.9]; Manse-CAP2b[Ala⁶]: A[2.0], E[1.0], F[1.0], L[0.9], R[1.0], Y[1.0], V[1.0]; Manse-CAP2b[Ala⁷]: A[2.1], E[0.7], F[1.0], L[1.1], P[1.1], Y[0.8], V[1.0]; Manse-CAP2b[Ala⁸]: A[2.0], E[1.0], F[1.0], L[1.0], P[0.9], R[1.0], Y[1.0]. The identities of the peptide analogs were confirmed via MALDI-TOF-MS on a Kratos Kompact Probe MALDI-TOF MS machine (Kratos Analytical, Ltd., Manchester, UK) with the presence of the following molecular ions (M + H⁺): Manse-CAP2b[2–8], 864.9 [M + H⁺]; Manse-CAP2b[3–8], 751.5 [M + H⁺]; Manse-CAP2b[4–8], 588.6 [M + H⁺]; Manse-CAP2b[5–8], 517.6 [M + H⁺]; Manse-CAP2b[Ala¹], 936.8 [M + H⁺]; Manse-CAP2b[Ala²], 933.8 [M + H⁺]; Manse-CAP2b[Ala³], 883.9 [M + H⁺]; Manse-CAP2b[Ala⁵], 899.5 [M + H⁺]; Manse-CAP2b[Ala⁶], 949.3 [M + H⁺]; Manse-CAP2b[Ala⁷], 890.6 [M + H⁺]; Manse-CAP2b[Ala⁸], 947.7 [M + H⁺].

2.3. Isolated housefly Malpighian tubule preparations

Fluid secretion from isolated housefly Malpighian tubules was measured using the “Ramsay assay” as previously described [10]. Tubules were removed from 3 to 4 day post-emergent adult female flies. Flies were dissected under *Musca* saline [10] and both anterior and posterior tubules were transferred to small (10 µl) drops of bathing fluid (a 1:1 mixture of saline and Schneider’s *Drosophila* medium) beneath water-saturated liquid paraffin in a Sylgard™ lined Petri dish. The tubules were allowed to equilibrate for 1 h before being challenged with test peptides. Urine escaped from the cut end of the tubule, which was pulled out into the liquid paraffin. Drops of urine were collected at 15 min intervals and their diameter (*d*) measured as they rested on the Sylgard base of the Petri dish using a Wild digital (MMS235) eyepiece micrometer. Urine volume was calculated as $\pi d^3/6$ and the rate of secretion obtained by dividing the secreted volume by the collection period. Data were normalized by expressing the increase in fluid secretion as a percentage of the response to 10 nM Musdo-K [16], which was added to all tubules at the end of each experiment. Dose-response curves were prepared using the computer program GraphPad Prism version 4.02 (GraphPad Software, San Diego, CA).

3. Results

A list of the fluid secretion activity of the two CAP2b/PVK peptides native to the housefly, Manse-CAP2b, Manse-CAP2b

Table 1 – Fluid secretion activity of CAP2b/PVK peptides and analogs on housefly (*M. domestica*) Malpighian tubules

Peptide	Sequence	Fluid secretion ^a	
		EC ₅₀ (nM) (95% CL)	Maximal response (%)
Musdo-CAP2b/PVK-1	AGGTSGLYAFPRVa	9 (7.2–11.1) [6]	100
Musdo-CAP2b/PVK-2	ASLFNAPRVa	102 (96–109) [6]	100
Manse-CAP2b	pELYAFPRVa	53 (39–72) [6]	
Manse-CAP2b[2–8]	LYAFPRVa	8 (3–22)	100
Manse-CAP2b[3–8]	YAFPRVa	428 (298–613)	100
Manse-CAP2b[4–8]	AFPRVa	Tr	
Manse-CAP2b[5–8]	FPRVa	Tr	
Manse-CAP2b[Ala ¹]	ALYAFPRVa	23 (11–46)	100
Manse-CAP2b[Ala ²]	pEAYAFPRVa	373 (255–544)	80
Manse-CAP2b[Ala ³]	pELAYAFPRVa	33 (10–105)	100
Manse-CAP2b[Ala ⁵]	pELYAAPRVa	115 (27–485)	70
Manse-CAP2b[Ala ⁶]	pELYAFARVa	89 (62–129)	100
Manse-CAP2b[Ala ⁷]	pELYAFPVa	Inactive	
Manse-CAP2b[Ala ⁸]	pELYAFPRAA	3500 (2250–5500)	45

^a Values are derived from dose–response curves based on data points derived from six replicates.

truncated analogs, and a Manse-CAP2b Ala scan series of analogs on housefly Malpighian tubules is presented in Table 1. The fluid secretion activity of *Manduca* CAP2b (EC₅₀ = 53 nM) lies midway between that of the native Musdo-CAP2b/PVK-1 (EC₅₀ = 9 nM) and Musdo-CAP2b/PVK-2 (EC₅₀ = 102 nM); and thus can serve as a reasonable model for a structure-activity relationship study of CAP2b/PVK peptides in the housefly. Interestingly, removal of the pE residue on the N-terminus of Manse-CAP2b leads to the C-terminal heptapeptide sequence common to both Manse-CAP2b and Musdo-CAP2b-1; and the diuretic activity of this fragment Manse-CAP2b[2–8] (EC₅₀ = 8 nM) matches that of Musdo-CAP2b-1 (EC₅₀ = 9 nM). The increase in activity observed with the loss of pGlu¹ likely arises from the fact that it now more closely matches the sequence of the native peptide. Removal of another amino acid residue from the N-terminus, as in Manse-CAP2b[3–8] leads to a large drop of an order of magnitude from the parent *Manduca* CAP2b peptide. Removals of a third (Manse-CAP2b[4–8]) and fourth (Manse-CAP2b[5–8]) residue lead to analogs that retain only trace activity. In the case of the Ala scan series, replacement of the Arg⁷ (Manse-CAP2b[Ala⁷]) led to an analog with no significant activity, indicating that Arg⁷ is therefore the most critical of residues. Replacement of Val⁸ (Manse-CAP2b[Ala⁸]; EC₅₀ = 3500 nM) led to a large drop in activity (70-fold) in comparison with the parent peptide, and demonstrated retention of only 45% of the maximal response of Manse-CAP2b and the native housefly peptides. Residues pGlu¹ (Manse-CAP2b[Ala¹]; EC₅₀ = 23 nM), Tyr³ (Manse-CAP2b[Ala³]; EC₅₀ = 33 nM), Phe⁵ (Manse-CAP2b[Ala⁵]; EC₅₀ = 115 nM), and Pro⁶ (Manse-CAP2b[Ala⁶]; EC₅₀ = 89 nM) would appear not to be critical for housefly diuretic activity as the resulting analogs were either a little more or a little less active. While Manse-CAP2b[Ala⁵] demonstrated a maximal response of 70%, this value is not statistically different from that of the parent peptide. Leu² would appear to be a semicritical residue, as Manse-CAP2b[Ala²] (EC₅₀ = 373 nM), proved to be about six-fold less active.

4. Discussion

The evaluation of the CAP2b sequence from *M. sexta*, Manse-CAP2b, in heterologous Malpighian tubule fluid secretion assays in fruit flies and houseflies [3,5,18] established that this family of peptides demonstrated diuretic activity in addition to cardioacceleratory activity. Seven of the eight residues of Manse-CAP2b are identical with the most potent of the native housefly CAP2b/PVK peptides, Musdo-CAP2b/PVK-1. As can be seen in Table 1, the EC₅₀ for housefly Malpighian tubule fluid secretion of Manse-CAP2b is midway between that of the two CAP2b/PVK peptides native to the housefly; and thus can serve as a relevant model sequence to determine a structure-activity relationship profile. The C-terminal fragment Manse-CAP2b[3–8] retains a significant portion of the Malpighian tubule fluid secretion activity of the parent peptide (Table 1). The active core, the minimum sequence required to retain significant diuretic activity in the housefly, is therefore the C-terminal hexapeptide. However, full activity requires the C-terminal heptapeptide fragment (Manse-CAP2b[2–8]), which is more active than the parent Manse-CAP2b and equipotent with the native Musdo-CAP2b-1 (Table 1). The results of the Ala-replacement series identify two residues, Arg⁷ and Val⁸, as critical for diuretic activity. The residue Leu² is semicritical; whereas all others are not critical. Not surprisingly, the critical (R⁷ and V⁸) and semicritical (L²) residues are among the most conserved among the CAP2b/PVK family throughout arthropods. The most conserved residues among the CAP2b/PVKs are the C-terminal PRVa-mide as well as a Leu at position 7 (equivalent to Leu² in Manse-CAP2b) from the C-terminus [27]. The structure-activity relationship profile determined in this study can aid in the design and development of peptidomimetic agonist/antagonist analogs of this diuretic peptide family with enhanced biostability and bioavailability as tools for arthropod endocrinologists and as potential pest management agents capable of disrupting the water balance in target insects.

Acknowledgments

This study was supported with a Collaborative Research Grant (#LST.CLG.979226) from the North Atlantic Treaty Organization (NATO) (RJN) and a Binational Agricultural Research and Development Grant (BARD #IS3356-02) (RJN), and a grant provided by the USDA/DOD DWFP Research Initiative (#0500-32000-001-01R) (RJN). In addition, we acknowledge the capable technical assistance of Allison Strey, Pawel Zubrzak and Nan Pryor of the USDA Areawide Pest Management Research Unit, Southern Plains Agricultural Research Center.

REFERENCES

- [1] Beyenbach KW. Transport mechanisms of diuresis in Malpighian tubules of insects. *J Exp Biol* 2003;206:3845–56.
- [2] Cazzamali G, Torp M, Hauser F, Williamson M, Grimmelikhuijzen CJP. The *Drosophila* gene CG9918 codes for a pyrokinin-1 receptor. *Biochem Biophys Res Commun* 2005;335:14–9.
- [3] Coast GM. The neuroendocrine regulation of salt and water balance in insects. *Zoology (ZACS)* 2001;103:179–88.
- [4] Coast GM, Orchard I, Phillips JE, Schooley DA. Insect diuretic and antidiuretic hormones. In: Evans PD, editor. *Advances in Insect Physiology*, London: Academic Press; 2002. p. 279–409.
- [5] Davies SA, Huesmann GR, Maddrell SH, O'Donnell MJ, Skaer NJV, Dow JAT, et al. CAP2b, a cardioacceleratory peptide, is present in *Drosophila* and stimulates fluid secretion via cGMP. *Am J Physiol* 1995;269:R1321–6.
- [6] Davis NT, Homberg U, Dirksen H, Levine RB, Hildebrand JG. Crustacean cardioactive peptide-immunoreactive neurons in the hawkmoth *Manduca sexta* and changes in their immunoreactivity during postembryonic development. *J Comp Neurol* 1993;338:612–27.
- [7] Eckert M, Herbert Z, Pollák E, Molnár L, Predel R. Identical cellular distribution of all abundant neuropeptides in the major abdominal neurohemal system of an insect (*Periplaneta americana*). *J Comp Neurol* 2002;452:264–75.
- [8] Eckert M, Predel R, Gundel M. Periviscerokinins-like immunoreactivity in the nervous system of the American cockroach. *Cell Tissue Res* 1999;295:159–70.
- [9] Gäde G, Auerswald L, Predel R, Marco HG. Substrate usage and its regulation during flight and swimming in the backswimmer, *Notonecta glauca*. *Physiol Entomol* 2004;29:84–93.
- [10] Holman GM, Nachman RJ, Coast GM. Isolation, characterization and biological activity of a diuretic myokinin neuropeptide from the housefly, *Musca domestica*. *Peptides* 1999;20:1–10.
- [11] Huesmann GR, Cheung CC, Loi PK, Lee TD, Swiderek K, Tublitz N. Amino acid sequence of CAP2b, an insect cardioacceleratory peptide from the tobacco hornworm *Manduca sexta*. *FEBS Lett* 1995;371:311–4.
- [12] Iversen A, Cazzamali G, Williamson M, Hauser F, Grimmelikhuijzen CJP. Molecular cloning and functional expression of a *Drosophila* receptor for the neuropeptides CAPA-1 and -2. *Biochem Biophys Res Commun* 2002;299:628–33.
- [13] Kean L, Cazenave W, Costes L, Boderick KE, Graham S, Pollock VP, et al. Two nitridergic peptides are encoded by the gene capability in *Drosophila melanogaster*. *Am J Physiol Regul Integr Comp Physiol* 2002;282:1297–307.
- [14] Loi PK, Tublitz NJ. Sequence and expression of the CAPA/CAP2b gene in the tobacco hawkmoth, *Manduca sexta*. *J Exp Biol* 2004;207:3681–91.
- [15] Meng X, Wahlström G, Immonen T, Kolmer M, Tirronen M, Predel R, et al. The *Drosophila hugin* gene codes for myostimulatory and ecdysis modifying neuropeptides. *Mech Dev* 2002;117:5–13.
- [16] Nachman RJ, Coast GM, Tichy SE, Russell DH, Miller JA, Predel R. Occurrence of insect kinins in the flesh fly, stable fly and horn fly—mass spectrometric identification from single nerves and diuretic activity. *Peptides* 2002;23:1885–94.
- [17] Nachman RJ, Isaac RE, Coast GM, Holman GM. Aib-containing analogues of the insect kinin neuropeptide family demonstrate resistance to an insect angiotensin-converting enzyme and potent diuretic activity. *Peptides* 1997;18:53–7.
- [18] Nachman RJ, Russell WK, Coast GM, Russell DH, Predel R. Mass spectrometric assignment of Leu/Ile in neuropeptides from single neurohemal organ preparations of insects. *Peptides* 2005;26:2151–6.
- [19] Nachman RJ, Russell WK, Coast GM, Russell DH, Miller JA, Predel R. Identification of PVK/CAP2b neuropeptides from single neurohemal organ preparations of the stable fly and horn fly via MALDI-TOF/TOF tandem mass spectrometry. *Peptides* 2006;27:521–6.
- [20] Nassel DR, Ohlsson LG, Cantera R. Metamorphosis of identified neurons innervating thoracic neurohemal organs in the blowfly—transformation of cholecystokinin like immunoreactive neurons. *J Comp Neurol* 1988;267:343–56.
- [21] Neupert S, Predel R, Russell WK, Davies R, Pietrantoni PV, Nachman RJ. Identification of tick periviscerokinins, the first neurohormone of Ixodidae: single cell analysis by means of MALDI-TOF/TOF mass spectrometry. *Biochem Biophys Res Commun* 2005;338:1860–4.
- [22] O'Brien MA, Taghert PH. A peritracheal neuropeptide system in insects: release of myomodulin-like peptides and ecdysis. *J Exp Biol* 1998;201:193–209.
- [23] Paluzzi J-P, Orchard I. Distribution, activity and evidence for the release of an anti-diuretic peptide in the kissing bug *Rhodnius prolixus*. *J Exp Biol* 2006;209:907–15.
- [24] Park Y, Kim YJ, Adams ME. Identification of G protein-coupled receptors for *Drosophila* PRXamide peptides, CCAP, corazonin, and AKH supports a theory of ligand-receptor coevolution. *Proc Natl Acad Sci USA* 2002;99:11423–8.
- [25] Predel R, Kellner R, Baggerman G, Steinmetzer T, Schoofs L. Identification of novel periviscerokinins from single neurohaemal release sites in insects MS/MS fragmentation complemented by Edman degradation. *Eur J Biochem* 2000;267:3869–74.
- [26] Predel R, Russell WK, Tichy SE, Russell DH, Nachman RJ. Capa-gene products in *Musca domestica* and *Neobellieria bullata*. *Peptides* 2003;24:1487–91.
- [27] Predel R, Wegener C. Periviscerokinins/CAPA gene related peptides in arthropods. *CMLS* 2006;63:2477–90.
- [28] Predel R, Wegener C, Russell WK, Tichy SE, Russell DH, Nachman RJ. Peptidomics of CNS-associated neurohemal systems of adult *Drosophila melanogaster*: a mass spectrometric survey of peptides from individual flies. *J Comp Neurol* 2004;474:379–92.
- [29] Quinlan MC, Tublitz NJ, O'Donnell MJ. Anti-diuresis in the blood-feeding insect *Rhodnius prolixus* Stål: the peptide CAP2b and cyclic GMP inhibit Malpighian tubule fluid secretion. *J Exp Biol* 1997;200:2363–7.
- [30] Santos JG, Pollak E, Rexer KH, Molnar L, Wegener C. Morphology and metamorphosis of the peptidergic Va neurons and the median nerve system of the fruitfly, *Drosophila melanogaster*. *Cell Tissue Res* 2006;326:187–99.

-
- [31] Sato Y, Oguchi M, Menjo N, Imai K, Saito H, Ikeda M, et al. Precursor polypeptide for multiple neuropeptides secreted from the subesophageal ganglion of the silkworm *Bombyx mori*: characterization of the cDNA encoding the diapause hormone precursor and identification of additional peptides. *Proc Natl Acad Sci USA* 1993;90:3251–5.
- [32] Wegener C, Reinl T, Jänsch L, Predel R. Direct mass spectrometric peptide profiling and fragmentation of larval peptide hormone release sites in *Drosophila melanogaster* reveals tagma-specific peptide expression and differential processing. *J Neurochem* 2006;96: 1362–74.